Covenant Medical Group • Orthopaedics

Orthopaedic Surgery & Sports Medicine • Garrett Kerns, DO

900 Cooper, Suite 3100 • Saginaw, Michigan 48602 • 989.583.7450 Tel • 989.583.7452 Fax

MEDICATION INSTRUCTIONS FOR DIRECT ANTERIOR TOTAL HIP ARTHROPLASTY

Cardiac Issues, Sulfa Allergy • 65 Years and Under

FIRST 24 HOURS PRIOR TO SURGERY:

- $\hfill\square$ Mobic® 15 mg once daily with food to be taken at 8:00 am.
- $\hfill\square$ OxyContin® 10 mg twice daily to be taken at 8:00 am and 8:00 pm.
- $\hfill\square$ Lyrica $^{\ensuremath{\oplus}}$ 50 mg twice daily to be taken at 8:00 am and 8:00 pm.

POST-OP MEDICATION REGIMEN:

- \Box Mobic 15 mg to be taken once daily with food until prescription is completed.
- □ Tylenol[®] 1000 mg to be taken 3 times daily for 1 week. Then can be take it as needed for pain. Patient needs to purchase over-the-counter Tylenol 500 mg tablets.
- □ Aspirin 81 mg to be taken twice daily for a total of 6 weeks for DVT prophylaxis.
- Dilaudid[®] 2 mg every 4 hours as needed for pain. 20 pills total.
- \Box Lyrica 50 mg twice daily to be taken at 8:00 am and 8:00 pm until the prescription is completed.



Extraordinary care for every generation.

©2018 Covenant HealthCare. All rights reserved. Bus. Dev. (AQ/RF) 2/18