

Covenant Medical Group • Orthopaedics

Orthopaedic Surgery & Sports Medicine • Garrett Kerns, DO

900 Cooper, Suite 3100 • Saginaw, Michigan 48602 • 989.583.7450 Tel • 989.583.7452 Fax

MEDICATION INSTRUCTIONS

FOR DIRECT ANTERIOR TOTAL HIP ARTHROPLASTY

High Risk DVT, Cardiac Issues, Sufa Allergy • 65 Years and Under

FIRST 24 HOURS PRIOR TO SURGERY:

- Mobic® 15 mg once daily with food to be taken at 8:00 am.
- OxyContin® 10 mg twice daily to be taken at 8:00 am and 8:00 pm.
- Lyrica® 50 mg twice daily to be taken at 8:00 am and 8:00 pm.

POST-OP MEDICATION REGIMEN:

- Mobic 15 mg to be taken once daily with food until prescription is completed.
- Tylenol® 1000 mg to be taken 3 times daily for 1 week. Then can be take it as needed for pain. Patient needs to purchase over-the-counter Tylenol 500 mg tablets.
- Lovenox® 40 mg subcutaneous injection once daily for a total of 4 weeks for DVT prophylaxis.
- Dilaudid® 2 mg every 4 hours as needed for pain. 20 pills total.
- Lyrica 50 mg twice daily to be taken at 8:00 am and 8:00 pm until the prescription is completed.

