

ACCELERATED REHABILITATION FOLLOWING ACL- PTG RECONSTRUCTION WITH LCL AND POSTEROLATERAL CORNER RECONSTRUCTION AND MEDIAL CAPSULAR INVOLVEMENT

I. PREOPERATIVE PHASE

Goals: Diminish inflammation, swelling, and pain
Restore normal range of motion (gradual knee extension)
Restore voluntary muscle activation
Provide patient education to prepare patient for surgery

Brace – Elastic wrap or knee sleeve to reduce swelling

Weight Bearing – As tolerated with or without crutches

Exercises:

- Ankle Pumps
- Passive knee extension (gradual progression)
- Passive knee flexion to tolerance
- Straight Leg Raises (3 Way, Flexion, Abduction), Pillow Squeezes
- Quadriceps Setting
- Closed kinetic chain exercises: mini squats, lunges, step-ups

Muscle Stimulation – Electrical muscle stimulation to quadriceps during voluntary quadriceps exercises (4 to 6 hours per day)

Neuromuscular/Proprioception Training -

- Eliminate quad avoidance gait
- Retro stepping drills
- Joint repositioning on Sports RAC
- Passive/active reposition at 90, 60, 30 degrees
- CKC squat/lunge repositioning on screen

Cryotherapy/Elevation – Apply ice 20 minutes of every hour, elevate leg with knee in full extension (knee must be above heart)

Patient Education – Review postoperative rehabilitation program
Review instructional video (optional)

Select appropriate surgical date

IMMEDIATE POST-OPERATIVE PHASE (Day 1 to Day 7)

Goals: Gradual passive knee extension
Diminish joint swelling and pain
Restore patellar mobility
Gradually improve knee flexion
Re-establish quadriceps control
Restore independent ambulation

Postoperative Day 1

Brace – EZ Wrap brace/Immobilizer applied to knee, locked in full extension during ambulation of Protonics

Weight Bearing – Two crutches, weight bearing as tolerated

Exercises:

- Ankle pumps
- Overpressure into passive knee extension
- Active and Passive knee flexion (90 degree by day 5)
- Straight leg raises (Flexion, Abduction), Pillow Squeezes
- Quadriceps isometric setting
- Hamstring stretches
- Closed kinetic chain exercises: mini squats, weight shifts

Muscle Stimulation – Use muscle stimulation during active muscle exercises (4-6 hours per day)

Continuous Passive Motion – As needed, 0 to 45/50 degrees (as tolerated and as directed by physician)

Ice and Evaluation – Ice 20 minutes out of every our and elevate with knee in full extension

Postoperative Day 2 to 3

Brace – EZ Wrap brace/Immobilizer, locked at zero degrees extension for ambulation and unlocked for sitting, etc.

Weight Bearing – Two crutches, weight bearing as tolerated

Range of Motion – Remove brace perform range of motion exercises 6-8 times per day
Perform frequent bouts of ROM to regain knee flexibility

Exercises:

- Multi-angle isometrics at 90 and 60 degrees (knee extension)
- Knee Extension 90-40 degrees
- Overpressure into extension (knee extension should be at least 0 degrees to slight hyperextension)
- Emphasize restoring knee extension
- Patellar mobilization
- Ankle pumps
- Straight leg raises, Pillow Squeezes
- Mini squats and weight shifts
- Quadriceps isometric setting

Muscle Stimulation – Electrical muscle stimulation to quads (6 hours per day)

Continuous Passive Motion – 0 to 90 degrees, as needed

Ice and Evaluation – Ice 20 minutes out of every hour and elevate leg with knee in full extension

Postoperative Day 4 to 7

Brace – EZ Wrap brace/Immobilizer, locked at zero degrees extension for ambulation and unlocked for sitting, etc.

Weight Bearing – Two Crutches weight bearing as tolerated

Range of Motion – Remove brace to perform range of motion exercises 6-8 times per day, knee flexion 90 degrees by day 5, approximately 100 degrees by day 7

Exercises:

- Multi-angle isometrics at 90 and 60 degrees (knee extension)
- Knee Extension 90-40 degrees
- Overpressure into extension
- Patellar mobilization (5-8 times daily)
- Ankle pumps
- Straight leg raises, Pillow Squeezes
- Mini squats and weight shifts
- Standing Hamstring curls
- Quadriceps isometric setting
- Proprioception and balance activities

Neuromuscular training/proprioception – OKC passive/active joint repositioning at 90, 60 degrees CKC squats/weight shifts with repositioning on sports RAC

Muscle Stimulation – Electrical muscle stimulation (continue 6 hours daily)

Continue Passive Motion – 0 to 90 degrees, as needed

Ice and Elevation – Ice 20 minutes of every hour and elevate leg with knee full extension

II. EARLY REHABILITATION PHASE (Week 2-4)

Criteria to Progress to Phase II

- 1) Quad Control (ability to perform good quad set and SLR)
- 2) Full passive knee extension
- 3) PROM 0-90 degrees
- 4) Good patellar mobility
- 5) Minimal joint effusion
- 6) Independent ambulation

Goals: Gradual increase to full passive knee extension
 Gradually increase knee flexion
 Diminish swelling and pain
 Muscle control and activation
 Restore proprioception/neuromuscular control
 Normalize patellar mobility

Week Two

Brace – Continue locked brace for ambulation

Weight Bearing – As tolerated (goal is to discontinue crutches 10-14 days post-op)

Passive Range of Motion – Self-ROM stretching (6-8 times daily), emphasis on maintaining full, passive range of motion

Exercises:

- Muscle stimulation to quadriceps exercises
- Isometric quadriceps sets
- Straight Leg raises (4 planes)
- Leg Press (0-60 degrees)
- Knee extension 90-40 degrees
- Half squats (0-40)
- Weight shifts
- Front and side lunges
- Uni-cam bicycle (low intensity cycling)
- Proprioception training
- Overpressure into extension
- Passive range of motion from 0 to 105 degrees
- Patellar mobilization
- Well leg exercises
- Progressive resistance extension program – start with 1 lb., progress 1 lb. per week

Proprioception/Neuromuscular Training

- OKC passive/active joint repositioning 90, 60, 30 degrees
- CKC joint repositioning during squats/lunges
- Initiate squats on tilt board use sports RAC with repositioning

Swelling control – Ice, compression, elevation

Week Three

Brace – Discontinue locked brace (some patients use ROM brace for ambulation)

Passive Range of Motion – Continue range of motion stretching and overpressure into extension (ROM should be 0-100/105 degrees)

Exercises:

- Continue all exercises as in week two
- Passive Range of Motion 0-105 degrees
- Bicycle for range of motion stimulus and endurance (emphasize ROM on bike)
- Pool walking program (if incision is closed)
- Eccentric quadriceps program 40-100 (isotonic only)
- Lateral lunges (straight plane)
- Front Step Downs
- Lateral Step-Overs (cones)
- Progress Proprioception drills, neuromuscular control drills
- Frequent bouts of ROM exercises

III. PROGRESSIVE STRENGTHENING/NEUROMUSCULAR CONTROL PHASE (Week 4-10)***Criteria to Enter Phase III***

- 1) Active Range of Motion 0-115 degrees
- 2) Quadriceps strength 60 % > contralateral side (isometric test at 60 degree knee flexion)
- 3) Unchanged KT Test bilateral values (+1 or less)
- 4) Minimal to no full joint effusion
- 5) No joint line or patellofemoral pain

Goals: Restore full knee range of motion (0 to 125 degrees)
Improve lower extremity strength
Enhance proprioception, balance, and neuromuscular control
Improve muscular endurance
Restore limb confidence and function

Brace – No immobilizer or brace, may use knee sleeve to control swelling/support

Range of Motion – Self-ROM (4-5 times daily using the other leg to provide ROM), emphasis on maintaining zero degrees passive extension
- PROM 0-125 degrees at 4 weeks

KT 2000 Test – (Week 4, 20 lb. anterior and posterior test)

Week 4

Exercises:

- Progress isometric strengthening program
- Leg Press (0-100 degrees)
- Knee extension 90 to 40 degrees
- Hip Abduction and Adduction
- Hip Flexion and Extension
- Lateral Step-Overs
- Lateral Lunges (straight plane and multi-plane drills)
- Lateral Step Ups
- Front Step Downs
- Wall Squats
- Vertical Squats
- Standing Toe Calf Raises
- Seated Toe Calf Raises
- Biodex Stability System (Balance, Squats, etc)
- Proprioception Drills
- Bicycle
- Stair Stepper Machine
- Pool Program (Backward Running, Hip and Leg Exercises)

Proprioception/Neuromuscular Drills:

- Tilt board squats (perturbation)
- Passive/active reposition OKC
- CKC repositioning on tilt board with sports RAC
- CKC lunges with sports RAC

Week 6

KT 2000 Test – 20 and 30 lb. anterior and posterior test

Exercises:

- Continue all exercises
- Pool running (forward) and agility drills
- Balance on tilt boards
- Progress to balance and ball throws
- Wall slides/squats

Week 8

KT 2000 Test – 20 and 30 lb. anterior and posterior test

Exercises:

- Continue all exercises listed in Weeks 4-6
- Leg Press Sets (single leg) 0-100 degrees and 40-100 degrees
- Plyometric Leg Press
- Perturbation Training
- Isokinetic exercises (90 to 40 degrees) (120 to 240 degrees/second)
- Walking Program
- Bicycle for endurance
- Stair Stepper Machine for endurance
- Biodex stability system

Week 10

KT 2000 Test – 20 and 30 lb. and Manual Maximum Test

Isokinetic Test – Concentric Knee Extension/Flexion at 180 and 300 degrees/second

Exercises:

- Continue all exercises listed in Weeks 6, 8 and 10
- Plyometric Training Drills
- Continue Stretching Drills
- Progress strengthening exercises and neuromuscular training

IV. ADVANCED ACTIVITY PHASE (Week 10-16)***Criteria to Enter Phase IV***

- 1) AROM 0-125 degrees or greater
- 2) Quad strength 75% of contralateral side, knee extension flexor:extensor ratio 70% to 75%
- 3) No change in KT values (Comparable with contralateral side, within 2 mm)
- 4) No pain or effusion
- 5) Satisfactory clinical exam
- 6) Satisfactory isokinetic test (values at 180 degrees)
 - Quadriceps bilateral comparison 75%
 - Hamstrings equal bilateral
 - Quadriceps peak torque/body weight 65% at 180o/s (males) 55% at 180o/s (females)
 - Hamstrings/quadriceps ratio 66% to 75%
- 7) Hop Test (80% of contralateral leg)
- 8) Subjective knee scoring (modified Noyes System) 80 points or better

Goals: Normalize lower extremity strength
 Enhance muscular power and endurance
 Improve neuromuscular control
 Perform selected sport-specific drills

Exercises:

- May initiate running program (weeks 10-12) if good quad control and ROM
- May initiate light sport program (golf)
- Continue all strengthening drills
 - Leg press
 - Wall squats
 - Hip Abd/Adduction

- Hip Flex/Ext
- Knee Extension 90-40
- Hamstring curls
- Standing toe calf
- Seated toe calf
- Step down
- Lateral step ups
- Lateral lunges
- Neuromuscular training
 - Lateral step-overs cones
 - Lateral lunges
 - Tilt board drills
 - Sports RAC repositioning on tilt board

Week 14-16

- Progress program
- Continue all drills above
- May initiate lateral agility drills
- Backward running

V. RETURN TO ACTIVITY PHASE (Week 16-22)

Criteria to Enter Phase V

- 1) Full Range of Motion
- 2) Unchanged KT 2000 Test (within 2.5 mm of opposite side)
- 3) Isokinetic Test that fulfills criteria
- 4) Quadriceps bilateral comparison (80% or greater)
- 5) Hamstring bilateral comparison (110% or greater)
- 6) Quadriceps torque/body weight ratio (55% or greater)
- 7) Hamstrings/Quadriceps ratio (70% or greater)
- 8) Proprioceptive Test (100% of contralateral leg)
- 9) Functional Test (85% or greater of contralateral side)
- 10) Satisfactory clinical exam
- 11) Subjective knee scoring (modified Noyes System) (90 points or better)

Goals: Gradual return to full-unrestricted sports
 Achieve maximal strength and endurance
 Normalize neuromuscular control
 Progress skill training

Tests – KT 2000, Isokinetic, and Functional Tests before return

Exercises

- Continue strengthening exercises
- Continue neuromuscular control drills
- Continue plyometrics drills
- Progress running and agility program
- Progress sport specific training
 - Running/cutting/agility drills
 - Gradual return to sport drills

6 MONTH FOLLOW-UP

Isokinetic test
KT 2000 test
Functional test

12 MONTH FOLLOW-UP

Isokinetic test
KT 2000 test
Functional test